

Organic Chemistry
Chemistry 210 and 211
Dr. Donald Robertson

Important Acid-Base Concepts for Organic Chemistry

Acids and bases are encountered in all areas of chemistry, sometimes even when we do not think we are using acids and bases. Knowing how to define acids and bases, as well as how to use them in chemical processes, is essential to understanding many organic chemistry reactions. The information contained here is designed to help you understand acids and bases better and how to use them in organic chemistry.

There are three basic definitions for acids and bases that have both historical and practical considerations. The first definition, described by Arrhenius, is used primarily for general chemistry. The second definition by Bronsted and Lowry applies most often to general and biochemistry, but has some applications for organic chemistry. The third definition by Lewis is applied most often to organic chemistry, but it is also applicable to all acids and bases, if you understand this definition fully. The Lewis definition of acids and bases is used primarily in non-aqueous (aprotic) solutions and is used to describe acids that do not contain ionizable hydrogens (non-proton donors).

The first useful definition of an acid was based on compounds that release H^+ ions. (An H^+ ion is often referred to as a proton, since a neutral hydrogen atom consists of one proton and one electron. When a hydrogen atom loses its lone electron, the only subatomic particle left is its proton; it is for this reason, therefore, that we call H^+ ions protons. In reality, the H^+ ion does not exist as a naked proton because it is always associated with an electronegative element, such as with the O in water to form H_3O^+ [oxonium or hydronium ion] or with the N in ammonia to produce NH_4^+ [ammonium ion] or their related derivatives.) This definition is referred to as the Arrhenius definition of acids. The common acids used in the laboratory are those that start their formula with an "H" atom (e.g., HCl, H_2SO_4 , HNO_3 , H_3PO_4 , etc.). In a similar manner, Arrhenius defined a base as a compound that releases OH^- (hydroxide) ions (e.g., NaOH, KOH, NH_4OH). The Arrhenius definition of acids and bases is very specific and applies only to compounds that meet these stringent requirements. The other definitions that follow are more general in nature.


Bronsted and Lowry introduced the second definition for acids and bases. Like Arrhenius, a Bronsted-Lowry acid is a proton (H^+ ion) donor. *Please note that the use of the word "proton" is actually part of this definition.* An acid (such as HCl), with a single ionizable hydrogen, is referred to as a monoprotic acid (i.e., it releases one proton). An acid (such as H_2SO_4), with two ionizable hydrogens, is referred to as a diprotic acid. An acid with three ionizable hydrogens (e.g., H_3PO_4), would be a triprotic acid, and so forth. According to Bronsted-Lowry, a base is a proton (H^+ ion) acceptor. There are two examples for bases that we will now consider. The first is the hydroxide ion (remember that the OH^- ion a base according to Arrhenius). The OH^- ion accepts (combines with) a proton (the H^+ ion), to form water. This is the typical acid-base reaction performed

during previous chemistry labs. However, please remember that the OH⁻ ion is not a base because it is the OH⁻ ion, per se.; it is a base, according to the Bronsted-Lowry definition, because it “accepts” a proton. The second example is that of ammonia (NH₃). Ammonia is a base, not because it contains the OH⁻ ion (which it obviously does not), but because it “accepts” a proton (H⁺ ion) to form the ammonium ion (NH₄⁺). The hallmark of Bronsted-Lowry bases is that they must possess at least one lone pair of electrons (you must be able to draw correct Lewis dot structures), and both water and ammonia have lone electron pairs (can you draw their correct Lewis dot structures?).

The third definition of an acid is that proposed by Lewis. These compounds are described as being Lewis acids or Lewis bases. You will eventually see that both the Arrhenius and Bronsted-Lowry definitions are consistent with those of Lewis, but you must be able to visualize correct Lewis dot structures. The Lewis definition of an acid is an electron pair acceptor (generally containing an atom with fewer than an octet of electrons; e.g., BH₃, AlCl₃, or a carbocation). You should be able to see that a proton (an H⁺ ion) meets this definition as well, since it can combine with a pair of electrons (e.g., donated by the OH⁻ ion) to react with the OH⁻ ion and form water. Conversely, a Lewis base is an electron pair donor (usually possessing an octet of electrons, but with at least one lone pair of electrons). So, how do you determine if you have a Lewis acid or base? You must be able to draw correct Lewis dot structures.

What are pK_a values, and how do you predict pK_a values?

The pK_a is a term that we encounter for acids that release protons. Let’s take the example of HCl(aq). When HCl(g) dissolves in water, it breaks apart into two parts, the proton (H⁺ ion) and its conjugate base, the chloride ion (Cl⁻). A conjugate base is formed when an acid loses a single proton. A conjugate acid is formed when a conjugate base accepts a proton. The ionization equation for HCl(g) dissolving in water is shown below:



Water also undergoes ionization according to the following equation:



The K_{eq} for water ionization is:

$$K_{\text{eq}} = [\text{H}^+][\text{OH}^-]/[\text{HOH}]$$

Which is the same equation for any equilibrium constant (K_{eq} equal the multiple of the concentrations of the products divided by multiple of the concentration of the reactants). For water, the concentration of H⁺ ions [H⁺] is 10⁻⁷ M, and the concentration of OH⁻ ions [OH⁻] is also 10⁻⁷ M; the concentration of water [HOH] is 55.555 M. When the concentrations of the various components are factored in, we get the following:

$$K_{\text{eq}} = [10^{-7}][10^{-7}]/[55.555] = 1.800018 \times 10^{-16}$$


The definition for the pK_a is the negative log of K_{eq} . Therefore,

$$pK_a = -\log [K_{eq}] = 15.7$$


which, if you recall, is the pK_a value that we have always used (and something you probably have already memorized) for water.

To determine the strength of an acid, the acid that has the smaller pK_a value (less positive, or more negative) is the stronger acid. For example, an acid with a pK_a of 1.0 is a stronger acid than an acid with a pK_a of 5.5; an acid with a pK_a of 15.7 is a stronger acid than an acid with pK_a of 36, etc. A pK_a value applies only to a compound or ion that has ionizable hydrogens (protons can be produced) on it. As a result, you need to identify compounds that have ionizable hydrogens. These include the common acids (HCl, HNO₃, H₂SO₄, HC₂H₃O₂; what are the names of these acids?). What about water, or ethanol, or ammonia; are these acids as well? As a consequence, a compound with an ionizable hydrogen is referred to as an acid (the Bronsted-Lowry definition). The conjugate base of an acid is formed when one, and *only* one, hydrogen comes off as a H⁺ ion (a proton).

Let's now look at sulfuric acid, H₂SO₄. When you remove one proton (H⁺), you produce its conjugate base, as shown below:



Please note that the conjugate base of sulfuric acid is not the sulfate ion. The reason is that it requires the removal of two protons to produce the sulfate ion from sulfuric acid, and a conjugate base can differ by only one proton from the acid. The hydrogen sulfate ion (HSO₄⁻) is actually the conjugate base of H₂SO₄. However, the hydrogen sulfate ion is also a conjugate acid, since it still has an ionizable hydrogen, which can be removed, as shown below:



Remember, in order to obtain a conjugate base, you look at the appropriate conjugate acid and remove one (*and only one*) H⁺ ion from it. *Be sure to remember that acid-base conjugate pairs differ by a single H⁺ ion.*

What is the formula for the conjugate base for each of the acids shown; what is the name of that conjugate base?

HCl Conjugate base is _____ base name _____

HOH Conjugate base is _____ base name _____

H₃PO₄ Conjugate base is _____ base name _____

H₂CO₃ Conjugate base is _____ base name _____

HCO₃⁻ Conjugate base is _____ base name _____

C₂H₅OH Conjugate base is _____ base name _____

Every acid, which has one or more ionizable hydrogens, has a conjugate base. Sometimes, the conjugate base may itself function as an acid (such as the hydrogen sulfate ion shown above), if it still has an ionizable hydrogen. Another example is the hydrogen carbonate ion (HCO₃⁻), also shown above.

At some time during your exposure to chemistry, you will encounter buffers. A buffer is a combination of conjugate acid and conjugate base that resists a change in pH. What is pH? It is, mathematically speaking, a value equal to the negative log of the [H⁺] concentration:



$$\text{pH} = -\log [\text{H}^+]$$

For example, if water has a [H⁺] concentration of 10⁻⁷ M, then the pH of water would be 7.0:

$$\text{pH} = -\log [\text{H}^+] = -\log [10^{-7}] = 7.0$$

An aqueous solution is neutral when the pH is 7.0, meaning that the concentration of [H⁺] and of [OH⁻] are equal 10⁻⁷ M. That is, if you have equal amounts of acid [H⁺] and of base [OH⁻], you are neither acidic nor basic, so as a result, the solution is neutral.

Now how does a buffer work? Let's assume we have the bicarbonate (hydrogen carbonate) ion: HCO₃⁻. If you examine this ion carefully, you will see that it has an ionizable hydrogen, hence it is an acid. Likewise, the hydrogen carbonate ion can accept a proton, so it is also a base. Look at what happens when the hydrogen carbonate ion reacts with both acid [H⁺] and base [OH⁻].



For a buffer, adding either acid or base does not change the pH (products are neither acidic or basic). In order to have a buffer, you must have an acid-base conjugate pair in solution.

Let's do this exercise with some other compounds, which we have already examined for relative acid strength above:

Predict the order (from most acidic to least acidic) of the following:



Most acidic: _____ > _____ > _____ > _____ Least acidic

Now, that you have ordered these compounds from most acidic (smallest pK_a) to least acidic (largest pK_a), how would you rank their conjugate bases?



Strongest base: _____ > _____ > _____ > _____ weakest base

How did you do? Remember that the strongest conjugate base comes from the weakest conjugate acid.

Nucleophiles and electrophiles

Organic chemistry is routinely broken down into acid-base reactions. Nucleophiles are nucleus-loving chemical species, meaning that a nucleophile is looking for an atom that is electron deficient (electrophile). For the most part, the electrophile is the acid, and the nucleophile is the base. According to the Lewis definition of acids and bases, a base (nucleophile) is an electron pair donor and a Lewis acid (electrophile) is an electron pair acceptor. Carbon atoms have been both electrophiles (e.g., carbocations and carbon atoms in S_N1 or S_N2 reactions). When we got to aromatic compounds, the carbon atoms functioned as nucleophiles, because the carbons in the aromatic ring structure had electrons available to them. However, no matter which class of carbon compounds you are working with, you will always need to identify what is the nucleophile and what is the electrophile for a particular reaction. This is not always easy, but there are a few clues. For example, when a carbon atom is bonded to an electronegative F, O, or N atom, that carbon will almost always be considered to be an electrophilic carbon. Therefore, it now becomes necessary to identify the compound or ion that is the nucleophile. A nucleophile will always have a free pair of electrons available on one of its atoms. This might take a little practice, but it will soon become obvious and easy to identify which compound or ion is the nucleophile and which compound or ion is the electrophile. Once you identify the nucleophile, all you need to do to get the reaction is to draw an arrow from the pair of electrons on the nucleophile to the electrophilic atom.

Most of the time, a carbon atom will be the electrophile. Let's just assume right now that it will be a carbon atom, and we need to identify which carbon will be the electrophilic carbon. Based on what we have learned before, any carbon attached to an O, N, or halogen (F, Cl, Br, I) will be the electrophile since it will have acquired a partial positive charge (+), due to differences in electronegativity. These compounds are rather easy to

identify and were encountered during substitution reactions. We will soon work with other types of carbon-containing compounds. The most common organic compounds that we will encounter will be those with carbonyl groups, which include the aldehydes, ketones, carboxylic acids, and the carboxylic acid derivatives such as esters, amides, acid anhydrides, and acid chlorides. The structures of these compounds will be examined later. However, what you need to know and look for in these compounds is that they all contain carbon atoms bonded to the electronegative element O. Because of the differences in electronegativity, the carbon atoms acquire a partial positive charge (δ^+), and are electrophilic. Do you understand this concept?

Now, since you have learned to identify the probable electrophile, you will need to know how to identify the probable nucleophile. In the past, the nucleophile was something like a halide (e.g., Br^- , Cl^- or I^-) or the OH^- ion. In addition, we also used the OH group of an alcohol as a nucleophile. In the future you will also need to identify certain carbon atoms that will function as a nucleophile (a carbon atom possessing an unbonded lone pair of electrons is referred to as a carbanion). We can wait until later to discuss these carbon nucleophiles, but the key concept is that we need to have a nucleophile and an electrophile for virtually every organic chemistry reaction we will be using.

Acids and bases, as electrophiles and nucleophiles, respectively, are encountered in virtually every organic chemistry reaction. Your success in organic chemistry will be directly proportional to your understanding of these concepts. Do not take them lightly, or you will struggle in the course. If you learn to understand these concepts, you will excel, not because you have memorized everything but because you have learned to think your way through the process. Try to look for unifying principles, and you will understand organic chemistry better than you ever imagined.

Good luck.